arXiv:hep-th/0108059v1 9 Aug 2001

TOPOLOGICAL FIELD THEORY APPROACH TO THE GENERALIZED BENNEY HIERARCHY

JEN-HSU CHANG[1] AND MING-HSIEN TU[2]
[1] *Department of Basic Course, Chung Cheng Institute of Technology,*
National Defense University,
Tashi, Taoyuan, Taiwan
E-mail: changjen@math.sinica.edu.tw
[2] *Department of Physics, National Chung Cheng University,*
Minghsiung, Chiayi, Taiwan
E-mail: phymhtu@ccunix.ccu.edu.tw

(February 1, 2008)

Abstract

The integrability of the generalized Benney hierarchy with three primary fields is investigated from the point of view of two-dimensional topological field theories coupled to gravity. The associated primary free energy and correlation functions at genus zero are obtained via Landau-Ginzburg formulation and the string equation is derived using the twistor construction for the Orlov operators. By adopting the approach of Dubrovin and Zhang we obtain the genus-one corrections of the Poisson brackets of the generalized Benney hierarchy.

Typeset using REVTEX

I. INTRODUCTION

Over the past ten years, the relationships between topological field theories (TFT) and integrable systems have attracted considerable interest [1]. It's Witten who realized that when the matter sectors of 2d TFT couple to gravity, the systems can be described by integrable hierarchies. In particular, Witten [2] and Kontsevich [3] showed that the partition function of the pure two-dimensional gravity, the simplest case, is equivalent to a particular tau function of the Korteweg-de Vries (KdV) hierarchy [4] characterized by the string equation. For nontrivial matter sectors such as A_n topological minimal models [5] and CP^1 topological σ-model [6–8] the integrable hierarchies behind them correspond to the dispersionless Kadomtsev-Petviashvili (dKP) and dispersionless Toda (dToda) hierarchies, respectively. Furthermore, the genus-zero primary free energy of a 2d TFT has been shown to satisfy the Witten-Dijkgraaf-Verlinde-Verlinde (WDVV) equation [2,9] of associativity which can be used to construct an integrable hierarchy of hydrodynamic type by taking into account the topological recursion relations of genus zero [2,10]. In fact it has been realized that the WDVV equation is intimately related to TFT and integrable hierarchies. On one hand, 2d TFT can be classified by the solutions of WDVV equations in the sense that a particular solution of WDVV provides the primary free energy of some topological model. On the other hand, for example, one can construct a particular tau function of the dispersionless Lax hierarchy, which, in small phase space, turned out to be a solution of WDVV equation and satisfies, in full phase space, the dispersionless Virasoro constraints [5] . Therefore the WDVV equation, being a bridge, enables us to study dispersionless integrable hierarchies from the point of view of 2d TFT and vice versa.

To formulate dispersionless Lax hierarchies (for a review, see [11]), let Λ be an algebra of Laurent series of the form

$$\Lambda = \{A | A = \sum_{i=-\infty}^{N} a_i p^i\},$$

with coefficients a_i depending on an infinite set of variables $T_1 \equiv X, T_2, T_3, \cdots$. Then one can define a Lie-bracket associated with Λ as follows [12]:

$$\{A, B\} = \frac{\partial A}{\partial p}\frac{\partial B}{\partial X} - \frac{\partial A}{\partial X}\frac{\partial B}{\partial p}, \qquad A, B \in \Lambda$$

which can be regarded as the Poisson bracket defined in the 2-dimensional phase space (X, p). The algebra Λ with respect to the above Poisson bracket can be decomposed into the Lie sub-algebras $\Lambda = \Lambda_{\geq k} \oplus \Lambda_{<k}$ for $k = 0, 1, 2$ where $\Lambda_{\geq k} = \{A \in \Lambda | A = \sum_{i \geq k} a_i p^i\}$, $\Lambda_{<k} = \{A \in \Lambda | A = \sum_{i<k} a_i p^i\}$ and we will use the notations : $\Lambda_+ = \Lambda_{\geq 0}$ and $\Lambda_- = \Lambda_{<0}$ for brevity. The Lax formulation of dispersionless hierarchies can be formally defined as

$$\frac{\partial \Lambda}{\partial T_n} = \{(\Lambda^{n/N})_{\geq k}, \Lambda\}, \quad n = 1, 2, 3, \cdots$$

where $\Lambda^{1/N}$ is the N-th root of Λ defined by $(\Lambda^{1/N})^N = \Lambda$.

- For $k = 0$, it's called dKP hierarchy [13] (see also [14]) and its relations to TFT have been investigated in [5,15,16].
- For $k = 1$, it's called dispersionless modified KP(dmKP) hierarchy [17–19] and its relations to TFT are under investigated [20]. It is the purpose of this paper.
- For $k = 2$, it's called dispersionless Harry-Dym hierarchy [17] and its relations to TFT are still open.

In this paper we continue our previous work [20] to study the Lax operator of the form [17,19]

$$L = p^2 + v^1 p + v^2 + v^3 p^{-1}, \tag{1.1}$$

which satisfies the non-standard Lax equations

$$\frac{\partial L}{\partial T_n} = \{L^{n/2}_{\geq 1}, L\}. \tag{1.2}$$

The first few nontrivial equations are

$$\frac{\partial}{\partial T_2}\begin{pmatrix} v^1 \\ v^2 \\ v^3 \end{pmatrix} = \begin{pmatrix} 2v^2_X \\ v^1 v^2_X + 2v^3_X \\ (v^1 v^3)_X \end{pmatrix},$$

$$8\frac{\partial}{\partial T_3}\begin{pmatrix} v^1 \\ v^2 \\ v^3 \end{pmatrix} = \begin{pmatrix} -3(v^1)^2 v^1_X + 12(v^2 v^1)_X + 24 v^3_X \\ 12 v^1_X v^3 + 24 v^1 v^3_X + 12 v^2 v^2_X + 3(v^1)^2 v^2_X \\ 12(v^2 v^3)_X + 3((v^1)^2 v^3)_X \end{pmatrix},$$

etc. They are multi-variable generalization of the Benney equation [21] which describes long wave in nonlinear phenomena and has very rich integrable structures [12,22–26]. Thus the set of the above equations is referred to as the generalized Benney hierarchy.

Recently, the bi-Hamiltonian structure associated with the generalized Benney hierarchy has been obtained from those structure of dmKP via truncation [17,19]. Starting from the bi-Hamiltonian structure the primary free energy of the corresponding 2d TFT can be constructed. The key idea relies on the fact that the Poisson brackets both are of hydrodynamic type introduced by Dubrovin and Novikov (DN) [27] and form a flat pencil (see below), in which, the geometric setting is provided by Frobenius manifolds [28–30]. One way to define such manifolds is to construct a function $F(t^1, t^2, \cdots, t^m)$ such that the associated functions,

$$c_{\alpha\beta\gamma} = \frac{\partial^3 F(t)}{\partial t^\alpha \partial t^\beta \partial t^\gamma}, \tag{1.3}$$

satisfy the following conditions [29]:

- The matrix $\eta_{\alpha\beta} = c_{1\alpha\beta}$ is constant and non-degenerate (for the discussion of degenerate cases, see [31]).

- The functions $c^\alpha_{\beta\gamma} = \eta^{\alpha\epsilon} c_{\epsilon\beta\gamma}$ define an associative and commutative algebra with a unity element. The associativity will give a system of non-linear partial differential equations for $F(t)$

  $$\frac{\partial^3 F(t)}{\partial t^\alpha \partial t^\beta \partial t^\lambda} \eta^{\lambda\mu} \frac{\partial^3 F(t)}{\partial t^\mu \partial t^\gamma \partial t^\sigma} = \frac{\partial^3 F(t)}{\partial t^\alpha \partial t^\gamma \partial t^\lambda} \eta^{\lambda\mu} \frac{\partial^3 F(t)}{\partial t^\mu \partial t^\beta \partial t^\sigma} \tag{1.4}$$

  which is the so-called WDVV equations [2,9] arising from TFT (see sec. 3).

- The functions F satisfies a quasi-homogeneity condition, namely, $\mathcal{L}_E F = d_F F + (\text{quadratic terms})$, where E is known as the Euler vector field.

On the other hand, given any solution(or primary free energy) of the WDVV equation, one can construct a Frobenius manifold $\mathcal{M}$ associated with it. On such a manifold one may interpret $\eta^{\alpha\beta}$ as a flat metric and t^α the flat coordinates. The associativity can be used to defines a Frobenius algebra on each tangent space $T^t\mathcal{M}$. Denoting the multiplication of the algebra as $u \cdot v$ then one may introduce a second flat metric on $\mathcal{M}$ defined by

$$g^{\alpha\beta} = E(dt^\alpha \cdot dt^\beta), \tag{1.5}$$

where $dt^\alpha \cdot dt^\beta = c^{\alpha\beta}_\gamma dt^\gamma = \eta^{\alpha\sigma} c^\beta_{\sigma\gamma} dt^\gamma$. This metric, together with the original metric $\eta^{\alpha\beta}$, define a flat pencil (i.e, $\eta^{\alpha\beta} + \lambda g^{\alpha\beta}$ is flat as well for any value of λ) which corresponds to the compatible condition in integrable systems and thus enables one to obtain a bi-Hamiltonian structure from a Frobenius manifold $\mathcal{M}$. The corresponding Hamiltonian densities are defined recursively by the formula [29]

$$\frac{\partial^2 h^{(n+2)}_\alpha}{\partial t^\beta \partial t^\gamma} = c^\sigma_{\beta\gamma} \frac{\partial h^{(n)}_\alpha}{\partial t^\sigma}, \qquad h^{(0)}_\alpha = \eta_{\alpha\beta} t^\beta \tag{1.6}$$

where the subscript $\alpha (= 1, 2, \cdots, m)$ labels the primary fields and the superscript $n \geq 0$, the level of gravitational descendants.

Our paper is organized as follows: In next section, we shall compute $c^\gamma_{\alpha\beta}$ from the generalized Benney hierarchy by using (1.6). We shall show that an additional hierarchy generated by Hamiltonians with logarithmic type emerges, which together with the ordinary Benney hierarchy are identified as renormalization group flows in genus-zero TFT coupled to 2d topological gravity. Then, in section 3, based on the fact that the dispersionless Lax operator (1.1) can be viewed as a superpotential in Landau-Ginzburg (LG) formulation of TFT [32,9], we define the fundamental correlation functions and derive their dynamic flows from the genus-zero topological recursion relations [2,10] of the associated TFT. In section 4, to establish the string equation describing the gravitational effect, we construct the twistor data for the generalized Benney hierarchy by using the Orlov operator of the dmKP hierarchy [18], as we have done in [20]. In section 5, we shall show that the Poisson brackets of the generalized Benney hierarchy can be unambiguously quantized by using the Dubrovin and Zhang (DZ) approach to bi-Hamiltonian structure in 2d TFT [33]. To do so, we compute the associated G-function, which is independent of t_1 and satisfies the following Getzler equations [34]

$$\sum_{1\leq\alpha_1,\alpha_2,\alpha_3,\alpha_4\leq 3} z_{\alpha_1}z_{\alpha_2}z_{\alpha_3}z_{\alpha_4}\left(3c^{\mu}_{\alpha_1\alpha_2}c^{\nu}_{\alpha_3\alpha_4}\frac{\partial^2 G}{\partial t^{\mu}\partial t^{\nu}} - 4c^{\mu}_{\alpha_1\alpha_2}c^{\nu}_{\alpha_3\mu}\frac{\partial^2 G}{\partial t^{\alpha_4}\partial t^{\nu}} - c^{\mu}_{\alpha_1\alpha_2}c^{\nu}_{\alpha_3\alpha_4\mu}\frac{\partial G}{\partial t^{\nu}}\right.$$
$$\left. +2c^{\mu}_{\alpha_1\alpha_2\alpha_3}c^{\nu}_{\alpha_4\mu}\frac{\partial G}{\partial t^{\nu}} + \frac{1}{6}c^{\mu}_{\alpha_1\alpha_2\alpha_3}c^{\nu}_{\alpha_4\mu\nu} + \frac{1}{24}c^{\mu}_{\alpha_1\alpha_2\alpha_3\alpha_4}c^{\nu}_{\mu\nu} - \frac{1}{4}c^{\mu}_{\alpha_1\alpha_2\nu}c^{\nu}_{\alpha_3\alpha_4\mu}\right) = 0, \qquad (1.7)$$

with $c^{\mu}_{\alpha_1\alpha_2\alpha_3\alpha_4} = \partial^2 c^{\mu}_{\alpha_1\alpha_2}/\partial t^{\alpha_3}\partial t^{\alpha_4}$, to evaluate the genus-one free energy $\mathcal{F}_1$. Section 6 is devoted to the concluding remarks.

II. INTEGRABILITY AND PRIMARY FREE ENERGY

In this section, we shall identify the primary free energy associated with the generalized Benney hierarchy from its integrability and construct the additional flows generated by the logarithmic Hamiltonians. According to the theory of Frobinus manifold, our starting point is the bi-Hamiltonian structure of the generalized Benney hierarchy, which can be constructed from those structure of the dmKP hierarchy via the Dirac reduction and thus the Lax flows (1.2) have the following bi-Hamiltonian description [17,19]:

$$\frac{\partial \mathbf{v}}{\partial T_n} = J_1\cdot\frac{\delta H_{n+2}}{\delta \mathbf{v}} = J_2\cdot\frac{\delta H_n}{\delta \mathbf{v}}$$

where $\mathbf{v}^T = (v^1, v^2, v^3)$, $(\delta H_i/\delta\mathbf{v})^T = (\delta H_i/\delta v^1, \delta H_i/\delta v^2, \delta H_i/\delta v^3)$ and

$$J_1 = \begin{pmatrix} 0 & 0 & 2\partial \\ 0 & 2\partial & v^1\partial \\ 2\partial & \partial v^1 & 0 \end{pmatrix}, \qquad J_2 = \begin{pmatrix} 6\partial & 4\partial v^1 & 2\partial v^2 \\ 4v^1\partial & v^2\partial + \partial v^2 + 2v^1\partial v^1 & 2\partial v^3 + v^3\partial + v^1\partial v^2 \\ 2v^2\partial & \partial v^3 + 2v^3\partial + v^2\partial v^1 & \partial v^1 v^3 + v^1 v^3\partial \end{pmatrix} \qquad (2.1)$$

with Hamiltonians defined by

$$H_n = \frac{2}{n}\int \mathrm{res}(L^{n/2})$$

where $\partial = \partial/\partial X$ and $\mathrm{res}(\Lambda)$ is the coefficient of p^{-1} of Λ. We list some of them:

$$\begin{aligned}
H_1 &= \int \left[v^2 - \frac{1}{4}(v^1)^2 \right], \\
H_2 &= \int v^3, \\
H_3 &= \int \left[\frac{1}{2} v^1 v^3 + \frac{1}{4}(v^2)^2 - \frac{1}{8} v^2 (v^1)^2 + \frac{1}{64}(v^1)^4 \right], \\
H_4 &= \int v^2 v^3, \\
H_5 &= \int \left[-\frac{1}{512}(v^1)^6 + \frac{3}{128}(v^1)^4 v^2 - \frac{1}{16}(v^1)^3 v^3 - \frac{3}{32}(v^1)^2 (v^2)^2 + \frac{3}{4} v^1 v^2 v^3 + \frac{1}{8}(v^2)^3 + \frac{3}{4}(v^3)^2 \right], \\
H_6 &= \int [v^1 (v^3)^2 + v^3 (v^2)^2].
\end{aligned}$$

To find the primary free energy associated with the generalized Benney hierarchy, we have to investigate the geometrical meaning of the Poisson brackets J_1 and J_2 which can be written as the DN type [27]

$$J_1^{ij} = \eta^{ij} \partial + \gamma_k^{ij} v_x^k, \qquad J_2^{ij} = g^{ij} \partial + \Gamma_k^{ij} v_x^k$$

with $J_1^{ij} = \partial J_2^{ij} / \partial v^2$ where γ_k^{ij} and Γ_k^{ij} are the contravariant Levi-Civita connections of the contravariant flat metrics η^{ij} and $g^{ij}(v)$ respectively. Let's introduce the flat coordinates

$$t^1 = v^2 - \frac{1}{4}(v^1)^2, \qquad t^2 = v^3, \qquad t^3 = v^1. \tag{2.2}$$

so that η^{ij} and g^{ij} can be expressed as follows:

$$\eta^{ij}(t) = \begin{pmatrix} 2 & 0 & 0 \\ 0 & 0 & 2 \\ 0 & 2 & 0 \end{pmatrix}, \qquad g^{ij}(t) = \begin{pmatrix} 2t^1 & 3t^2 & t^3 \\ 3t^2 & 2t^2 t^3 & 2t^1 + \frac{1}{2}(t^3)^2 \\ t^3 & 2t^1 + \frac{1}{2}(t^3)^2 & 6 \end{pmatrix}, \tag{2.3}$$

with $\partial g^{ij} / \partial t^1 = \eta^{ij}$. It is easy to show that t^α are just the densities of the Casimirs invariants for the first structure J_1.

Now we are in a position to compute the structure coefficients $c_{\beta\gamma}^{\sigma}$ of the generalized Benney hierarchy. Choosing $h_1^{(0)} = \eta_{1\beta} t^\beta = t^1/2$ and $h_3^{(0)} = \eta_{3\beta} t^\beta = t^2/2$, the equations (1.6) implies

$$h_{2n-1} = \frac{2^{n-1}}{(2n-1)!!} \mathrm{res} L^{n-1/2}, \qquad h_1 = h_1^{(0)} = \frac{t^1}{2}, \tag{2.4}$$

$$h_{2n} = \frac{2^{n-1}}{(2n)!!} \mathrm{res} L^n, \qquad h_2 = h_3^{(0)} = \frac{t^2}{2}. \tag{2.5}$$

with non-zero coefficients

$$\begin{aligned}
&c^1_{11} = 1, \quad c^1_{23} = 1, \\
&c^2_{12} = 1, \quad c^2_{23} = \frac{t^3}{2}, \quad c^2_{33} = \frac{t^2}{2}, \\
&c^3_{13} = 1, \quad c^3_{22} = \frac{2}{t^2}, \quad c^3_{33} = \frac{t^3}{2}.
\end{aligned}$$

Then from (1.3), we immediately obtain the free energy

$$F(t^1, t^2, t^3) = \frac{1}{12}(t^1)^3 + \frac{1}{2}t^1t^2t^3 + \frac{1}{24}(t^3)^3t^2 + \frac{1}{2}(t^2)^2\left(\log t^2 - \frac{3}{2}\right)$$

which is indeed a finite-dimensional solution to the WDVV equations (1.4). Also, from (1.5) and (2.3), we have the associated Euler vector field

$$E = t^1\frac{\partial}{\partial t^1} + \frac{3}{2}t^2\frac{\partial}{\partial t^2} + \frac{1}{2}t^3\frac{\partial}{\partial t^3},$$

which implies the quasi-homogeneity condition: $\mathcal{L}_E F(t) = 3F(t) + \frac{3}{4}(t^2)^2$.

Next we turn to the hierarchy corresponding to $\alpha = 2$ in (1.6) with $h_2^{(0)} = \eta_{2\beta}t^\beta = t^3/2$ and obtain

$$\begin{aligned}
\bar{h}_0 &= \frac{t^3}{2}, \\
\bar{h}_1 &= \frac{(t^3)^3}{24} + \frac{1}{2}t^1t^3 + t^2\left(\log t^2 - \frac{5}{2}\right), \\
\bar{h}_2 &= \frac{(t^3)^5}{320} + \frac{t^1(t^3)^3}{24} + \frac{(t^1)^2t^3}{4} + \frac{1}{4}(t^3)^2t^2\left(\log t^2 - \frac{3}{2}\right) + t^1t^2\left(\log t^2 - \frac{5}{2}\right),
\end{aligned}$$

As in the work [20], the Hamiltonian densities $h_2^{(n)} = \bar{h}_n$ can be expressed as

$$\bar{h}_n = \frac{3}{2n!}\mathrm{res}[L^n(\log L - c_n)], \tag{2.6}$$

where we use the following prescription for $\log L$

$$\begin{aligned}
\log L &= \log(p^2 + v^1p + v^2 + v^3p^{-1}) \\
&= \frac{1}{3}\log[p^2(1 + v^1p^{-1} + v^2p^{-2} + v^3p^{-3})] + \frac{2}{3}\log\left[v^3p^{-1}\left(1 + \frac{v^1}{v^3}p + \frac{v^2}{v^3}p^2 + \frac{1}{v^3}p^3\right)\right] \\
&= \frac{2}{3}\log v^3 + \frac{1}{3}\log(1 + v^1p^{-1} + v^2p^{-2} + v^3p^{-3}) + \frac{2}{3}\log\left(1 + \frac{v^1}{v^3}p + \frac{v^2}{v^3}p^2 + \frac{1}{v^3}p^3\right),
\end{aligned} \tag{2.7}$$

and $c_n - c_{n-1} = 1/n,\ c_0 = 1$. The Lax and bi-Hamiltonian formulations corresponding to $\bar{T}_n$-flows can be summarized as

$$\begin{aligned}\frac{\partial L}{\partial \bar{T}_n} &= 2\{\bar{B}_n, L\}, \qquad \bar{B}_n = [L^n(\log L - c_n)]_{\geq 1} \\ &= \{\bar{H}_{n+1}, L\}_1 = \{\bar{H}_n, L\}_2, \qquad \bar{H}_n = 2(n-1)! \int \bar{h}_n \end{aligned} \tag{2.8}$$

Notice that the additional flows generated by the Hamiltonians $\bar{H}_n$ are compatible with those flows of the ordinary Benney hierarchy (1.2). We will see later that the forms (2.4) , (2.5) and (2.6) can be used to formulate the generalized Benney hierarchy as a 2d TFT coupled to gravity via LG description.

Before ending this section we would like to remark that the second Poisson bracket J_2 in fact reveals the classical realization of w_3-$U(1)$-Kac-Moody algebra [19] with t^1 being the Diff S^1 tensor of weight 2, and t^2 and t^3, tensors of weights 3 and 1, respectively. This is due to the fact that the Diff S^1 flows are just the Hamiltonian flows generated by the Hamiltonian $H_1 = \int t^1/2$.

III. LANDAU-GINZBURG FORMULATION

In this section we shall set up the correspondence between the Benney hierarchy and its associated TFT at genus zero. Let us first briefly recall some basic notions in TFT for convenience.

A topological matter theory can be characterized by a set of BRST invariant observables $\{\mathcal{O}_1, \mathcal{O}_2, ...\}$ with couplings $\{T^\alpha\}$ where $\mathcal{O}_1$ denotes the identity operator. If the number of observables is finite the theory is called topological minimal model and the observables are referred to the primary fields. When the theory couples to gravity, a set of new observables emerge as gravitational descendants $\{\sigma_n(\mathcal{O}_\alpha), n = 1, 2, \cdots\}$ with new coupling constants $\{T^{\alpha,n}\}$. The identity operator $\mathcal{O}_1$ now becomes the puncture operator P. For convenience we shall identify the primary fields $\mathcal{O}_\alpha$ and the coupling constants T^α to $\sigma_0(\mathcal{O}_\alpha)$ and $T^{\alpha,0}$, respectively. As usual, we shall call the space spanned by $\{T^{\alpha,n}, n = 0, 1, 2, \cdots\}$ the full phase space and the subspace parametrized by $\{T^\alpha\}$ the small phase space. Hence, for a topological minimal model, the small phase space is finite-dimensional.

For a topological model the most important quantities are correlation functions which do not depend on coordinates and describe the underlying topological properties of the manifold. The generating function of correlation functions is the full free energy defined by

$$\mathcal{F}(T) = \sum_{g=0}^{\infty} \mathcal{F}_g(T) = \sum_{g=0}^{\infty} \langle e^{\sum_{\alpha,n} T^{\alpha,n}\sigma_n(\mathcal{O}_\alpha)} \rangle_g$$

where $\langle \cdots \rangle_g$ denotes the expectation value on a Riemann surface of genus g with respect to a classical action. In the subsequent sections, we will omit the exponential factor without causing any confusion. Therefore a generic m-point correlation function can be calculated as follows

$$\langle \sigma_{n_1}(\mathcal{O}_{\alpha_1})\sigma_{n_2}(\mathcal{O}_{\alpha_2})\cdots\sigma_{n_m}(\mathcal{O}_{\alpha_m})\rangle_g = \frac{\partial^m \mathcal{F}_g}{\partial T^{\alpha_1,n_1}\partial T^{\alpha_2,n_2}\cdots\partial T^{\alpha_m,n_m}}.$$

In the following, we shall restrict ourselves to the trivial topology, i.e., the genus-zero sector ($g = 0$) since this part is more relevant to dispersionless integrable hierarchies. In particular, the genus-zero free energy restricting on the small phase space is the primary free energy defined by

$$\mathcal{F}_0|_{T^\alpha = t^\alpha, T^{\alpha,n\geq 1}=0} = F(t).$$

Let us consider some genus-zero correlation functions on the small phase space. The metric on the space of primary fields is defined by

$$\langle P\mathcal{O}_\alpha\mathcal{O}_\beta\rangle = \eta_{\alpha\beta}.$$

When $\eta_{\alpha\beta}$ is independent of the couplings we call it the flat metric and the couplings T^α the flat coordinates. In fact, the three-point functions in the small phase space can be expressed as

$$\langle \mathcal{O}_\alpha\mathcal{O}_\beta\mathcal{O}_\gamma\rangle = \frac{\partial F}{\partial T^\alpha \partial T^\beta \partial T^\gamma} \equiv c_{\alpha\beta\gamma}$$

which provide the structure constants of the commutative and associative algebra $\mathcal{O}_\alpha\mathcal{O}_\beta = c^\gamma_{\alpha\beta}\mathcal{O}_\gamma$. The associativity of $c^\gamma_{\alpha\beta}$, i.e., $c^\mu_{\alpha\beta}c^\sigma_{\mu\gamma} = c^\mu_{\alpha\gamma}c^\sigma_{\mu\beta}$, will give the WDVV equations (1.4).

Since the Lax hierarchy under consideration is a three-variable theory, thus only three primary fields $\{\mathcal{O}_1 = \mathcal{O}_P \equiv P, \mathcal{O}_2 = \mathcal{O}_Q \equiv Q, \mathcal{O}_3 = \mathcal{O}_R \equiv R\}$ are involved in the TFT formulation and we shall identify

$$v^1|_{T^{\alpha,n\geq 1}=0} = T^R, \qquad v^2|_{T^{\alpha,n\geq 1}=0} = T^P + \frac{1}{4}(T^R)^2, \qquad v^3|_{T^{\alpha,n\geq 1}=0} = T^Q$$

on the small phase space and the Lax operator in small phase space is written as

$$L(z) = z^2 + T^R z + T^P + \frac{1}{4}(T^R)^2 + T^Q z^{-1}$$

which can be viewed as a rational superpotential in LG formulation of TFT [32,9]. According to the LG theory, the primary fields corresponding to T^α are defined by

$$\mathcal{O}_P(z) = \frac{\partial L(z)}{\partial T^P} = 1, \qquad \mathcal{O}_Q(z) = \frac{\partial L(z)}{\partial T^Q} = z^{-1}, \qquad \mathcal{O}_R(z) = \frac{\partial L(z)}{\partial T^Q} = z + \frac{T^R}{2}$$

which can be used to compute the three-point correlation functions through the residue formula [32,9]:

$$c_{\alpha\beta\gamma} = \mathrm{Res}_{L'=0}\left[\frac{\mathcal{O}_\alpha(z)\mathcal{O}_\beta(z)\mathcal{O}_\gamma(z)}{\partial_z L(z)}\right]. \tag{3.1}$$

It is easy to show that (3.1) reproduces the previous $c_{\alpha\beta\gamma}$ and F on the small phase space. In particular, the non-zero components of the flat metric on the space of primary fields are given by $\eta_{PP} = \eta_{QR} = \eta_{RQ} = 1/2$, $\eta^{PP} = \eta^{QR} = \eta^{RQ} = 2$ as we obtained previously. Now we define the fundamental correlation functions $\langle \mathcal{O}_\alpha \mathcal{O}_\beta \rangle = \partial^2 F/\partial T^\alpha \partial T^\beta$ as follows:

$$\begin{aligned}
\langle PP \rangle = \frac{T^P}{2}, \quad \langle PQ \rangle = \frac{T^R}{2}, \quad \langle PR \rangle = \frac{T^Q}{2}, \\
\langle QQ \rangle = \log T^Q, \quad \langle QR \rangle = \frac{T^P}{2} + \frac{(T^R)^2}{8}, \quad \langle RR \rangle = \frac{T^Q T^R}{4},
\end{aligned}$$

Although these two-point correlation functions are defined on the small phase space, however, it has been shown [10] that they can be defined on the full phase space through the variables v^1, v^2 and v^3 in which the gravitational couplings $T^{\alpha,n}$ do not vanish. Hence one can write down these genus-zero two-point functions on the full phase space and obtain the following constitutive relations:

$$\begin{aligned}
\langle PP \rangle = \frac{v^2}{2} - \frac{(v^1)^2}{8}, \quad \langle PQ \rangle = \frac{v^1}{2}, \quad \langle PR \rangle = \frac{v^3}{2}, \\
\langle QQ \rangle = \log v^3, \quad \langle QR \rangle = \frac{v^2}{2}, \quad \langle RR \rangle = \frac{v^1 v^3}{4},
\end{aligned} \tag{3.2}$$

which will be important to provide the connection between the generalized Benney hierarchy and its associated TFT.

Based on the constitutive relations (3.2), we can identify the gravitational flows for v^α in the full phase space as the integrable flows by taking into account the genus-zero topological recursion relation [2,10]:

$$\langle \sigma_n(\mathcal{O}_\alpha) AB \rangle = \sum_{\beta,\gamma=P,Q,R} n \langle \sigma_{n-1}(\mathcal{O}_\alpha) \mathcal{O}_\beta \rangle \eta^{\beta\gamma} \langle \mathcal{O}_\gamma AB \rangle, \qquad \alpha = P, Q, R. \tag{3.3}$$

For example, setting $n = 1$, $\mathcal{O}_\alpha = P$ and $A = P, B = Q$ then

$$\begin{aligned}
\frac{\partial v^1}{\partial T^{P,1}} &= 2\langle \sigma_1(P) PQ \rangle \\
&= 4[\langle PP \rangle \langle PPQ \rangle + \langle PQ \rangle \langle RPQ \rangle + \langle PR \rangle \langle QPQ \rangle \\
&= 4[\langle PP \rangle \langle PQ \rangle' + \langle PQ \rangle \langle RQ \rangle' + \langle PR \rangle \langle QQ \rangle'] \\
&= \left[v^1 v^2 - \frac{1}{12}(v^1)^3 + 2v^3 \right]_X
\end{aligned}$$

where we denote $f' = \partial f/\partial T^P = \partial f/\partial X$. Similarly, taking $\mathcal{O}_\alpha = R$ we have

$$\frac{\partial v^1}{\partial T^{R,1}} = 2\langle \sigma_1(R) PQ \rangle = \left[v^1 v^3 + \frac{1}{2}(v^2)^2 \right]_X .$$

On the other hand, taking $n = 2$ we get

$$
\begin{aligned}
\frac{\partial v^1}{\partial T^{P,2}} &= 2\langle \sigma_2(P) PQ \rangle \\
&= \left[\frac{1}{80}(v^1)^5 - \frac{1}{6}(v^1)^3 v^2 + (v^1)^2 v^3 + v^1 (v^2)^2 + 4v^2 v^3 \right]_X , \\
\frac{\partial v^1}{\partial T^{R,2}} &= 2\langle \sigma_2(R) PQ \rangle \\
&= \left[\frac{1}{3}(v^2)^3 + 2v^1 v^2 v^3 + (v^3)^2 \right]_X .
\end{aligned}
$$

etc. We present other flows in Appendix A.

Next we turn to the $T^{Q,n}$-flows. Choosing $\mathcal{O}_\alpha = Q$ and using the topological recursion relation (3.3), we obtain

$$
\begin{aligned}
\frac{\partial v^1}{\partial T^{Q,1}} &= \left[\frac{1}{2}(v^1)^2 + 2v^2 \log v^3 \right]_X , \\
\frac{\partial v^1}{\partial T^{Q,2}} &= \left[-\frac{1}{12}(v^1)^4 + v^2 (v^1)^2 + 2(v^2)^2 \log v^3 + 4v^1 v^3 (\log v^3 - 1) \right]_X .
\end{aligned}
$$

Comparing with the Lax equations (1.2) and (2.8) it turns out that the above recursion relations can be recasted into the Lax form in terms of coupling times $T^{\alpha,n}$ as

$$
\begin{aligned}
\frac{\partial L}{\partial T^{P,n}} &= \frac{2^n n!}{(2n+1)!!} \{ L_{\geq 1}^{n+1/2}, L \}, \\
\frac{\partial L}{\partial T^{Q,n}} &= \{ [L^n (\log L - c_n)]_{\geq 1}, L \}. \\
\frac{\partial L}{\partial T^{R,n}} &= \frac{1}{2n+2} \{ L_{\geq 1}^{n+1}, L \}
\end{aligned} \tag{3.4}
$$

with the identification

$$
T_{2n-1} = \frac{2^{n-1}(n-1)!}{(2n-1)!!} T^{P,n-1}, \quad T_{2n} = \frac{1}{2n} T^{R,n-1}, \quad \bar{T}_n = \frac{1}{2} T^{Q,n}. \tag{3.5}
$$

On the other hand, the associated commuting Hamiltonian flows with respect to the bi-Hamiltonian structures are thus given by

$$
\begin{aligned}
\frac{\partial \mathbf{v}}{\partial T^{P,n}} &= \{ H_{P,n+1}, \mathbf{v} \}_1 = \frac{2n}{2n+1} \{ H_{P,n}, \mathbf{v} \}_2, \qquad H_{P,n} = \frac{2^n (n-1)!}{(2n+1)!!} \int \mathrm{res} L^{n+1/2} \\
\frac{\partial \mathbf{v}}{\partial T^{R,n}} &= \{ H_{R,n+1}, \mathbf{v} \}_1 = \frac{n}{n+1} \{ H_{R,n}, \mathbf{v} \}_2, \qquad H_{R,n} = \frac{1}{2n(n+1)} \int \mathrm{res} L^{n+1} \\
\frac{\partial \mathbf{v}}{\partial T^{Q,n}} &= \{ H_{Q,n+1}, \mathbf{v} \}_1 = \{ H_{Q,n}, \mathbf{v} \}_2, \qquad H_{Q,n} = \frac{3}{2n} \int \mathrm{res}[L^n (\log L - c_n)]
\end{aligned} \tag{3.6}
$$

where $n = 0, 1, 2, \cdots$.

Furthermore, using (2.2), (3.2) and (3.4) the flows for t^α are

$$\begin{aligned}
\frac{\partial t^1}{\partial T^{\alpha,n}} &= 2\langle\sigma_n(\mathcal{O}_\alpha)PP\rangle = (R^{(1)}_{\alpha,n})', \\
\frac{\partial t^2}{\partial T^{\alpha,n}} &= 2\langle\sigma_n(\mathcal{O}_\alpha)PR\rangle = (R^{(2)}_{\alpha,n})', \\
\frac{\partial t^3}{\partial T^{\alpha,n}} &= 2\langle\sigma_n(\mathcal{O}_\alpha)PQ\rangle = (R^{(3)}_{\alpha,n})'
\end{aligned} \tag{3.7}$$

where $R^{(\beta)}_{\alpha,n}$ are the analogues of the Gelfand-Dickey potentials [4] of the KP hierarchy, given by

$$\begin{aligned}
R^{(1)}_{P,n} &= 2\langle\sigma_n(P)P\rangle = \frac{2^{n+1}n!}{(2n+1)!!}\mathrm{res}(L^{n+1/2}), \\
R^{(1)}_{Q,n} &= 2\langle\sigma_n(Q)P\rangle = 2\mathrm{res}[L^n(\log L - c_n)], \\
R^{(1)}_{R,n} &= 2\langle\sigma_n(R)P\rangle = \frac{1}{n+1}\mathrm{res}(L^{n+1}), \\
R^{(2)}_{P,n} &= 2\langle\sigma_n(P)R\rangle = \frac{2^n n!}{(2n+1)!!}[2(L^{n+1/2})_{-2} + v^1\mathrm{res}(L^{n+1/2})], \\
R^{(2)}_{Q,n} &= 2\langle\sigma_n(Q)R\rangle = [2(L^n(\log L - c_n))_{-2} + v^1\mathrm{res}(L^n(\log L - c_n))], \\
R^{(2)}_{R,n} &= 2\langle\sigma_n(R)R\rangle = \frac{1}{2n+2}[2(L^{n+1})_{-2} + v_1\mathrm{res}(L^{n+1})], \\
R^{(3)}_{P,n} &= 2\langle\sigma_n(P)Q\rangle = \frac{2^{n+1}n!}{(2n+1)!!}(L^{n+1/2})_0, \\
R^{(3)}_{Q,n} &= 2\langle\sigma_n(Q)Q\rangle = 2[L^n(\log L - c_n)]_0, \\
R^{(3)}_{R,n} &= 2\langle\sigma_n(R)Q\rangle = \frac{1}{n+1}(L^{n+1})_0.
\end{aligned} \tag{3.8}$$

In particular, the two-point correlators involving the first and the second descendants are also presented in Appendix A.

Finally, we would like to remark that the first three equations of (3.8) can be integrated to yield

$$\begin{aligned}
\langle\sigma_n(P)\rangle &= \frac{2^{n+1}n!}{(2n+3)!!}\mathrm{res}L^{n+3/2}, \\
\langle\sigma_n(Q)\rangle &= \frac{1}{n+1}\mathrm{res}[L^{n+1}(\log L - c_{n+1})], \\
\langle\sigma_n(R)\rangle &= \frac{1}{2(n+1)(n+2)}\mathrm{res}L^{n+2}
\end{aligned}$$

which represent the one-point functions of gravitational descendants at genus zero in the LG description.

IV. TWISTOR CONSTRUCTION FOR THE STRING EQUATION

After coupling to gravity, the dynamical variables v^α (or flat coordinates t^α) become functions on the full phase space that is parametrized by $T^{\alpha,n}$. Then v^α (or t^α) satisfy not only the Lax flows but also the string equation which characterizes the corresponding additional symmetries or W-constraint associated with the generalized Benney hierarchy. Since the Lax formulation in sec. 3 for the generalized Benney hierarchy is similar to that of the dToda type hierarchy [14,11]. This motivates us to reproduce the generalized Benney equations by imposing constraints on the Lax operators and the associated Orlov operators of the dmKP hierarchy [18] through the twistor data. We shall follow closely that of [35] to show that the constraints imposing on the twistor data implies the string equation of the generalized Benney hierarchy. Let us consider two Lax operators μ and $\tilde{\mu}$ with the following Laurent expansions ($T^1 = X$):

$$\mu = p + \sum_{n=0}^{\infty} v^n(T,\tilde{T})p^{-n}, \qquad \tilde{\mu}^{-1} = \tilde{v}_0(T,\tilde{T})p^{-1} + \sum_{n=0}^{\infty} \tilde{v}_{n+1}(T,\tilde{T})p^n \tag{4.1}$$

which satisfy the commuting Lax flows

$$\begin{aligned} \frac{\partial\mu}{\partial T_n} &= \{B_n, \mu\}, \qquad & \frac{\partial\mu}{\partial \tilde{T}_n} &= \{\tilde{B}_n, \mu\}, \\ \frac{\partial\tilde{\mu}}{\partial T_n} &= \{B_n, \tilde{\mu}\}, \qquad & \frac{\partial\tilde{\mu}}{\partial \tilde{T}_n} &= \{\tilde{B}_n, \tilde{\mu}\}, \end{aligned} \tag{4.2}$$

where

$$B_n \equiv (\mu^n)_{\geq 1}, \qquad \tilde{B}_n \equiv (\tilde{\mu}^{-n})_{\leq 0}, \quad (n = 1,2,3\cdots)$$

Next, we consider the Orlov operators corresponding to dmKP

$$M = \sum_{n=1}^{\infty} nT_n\mu^{n-1} + \sum_{n=1}^{\infty} w_n(T,\tilde{T})\mu^{-n}, \qquad \tilde{M} = -\sum_{n=1}^{\infty} n\tilde{T}_n\tilde{\mu}^{-n-1} + X + \sum_{n=1}^{\infty} \tilde{w}_n(T,\tilde{T})\tilde{\mu}^{n} \tag{4.3}$$

with constraint

$$\{\mu, M\} = 1, \qquad \{\tilde{\mu}, \tilde{M}\} = 1. \tag{4.4}$$

In fact, the coefficient functions w_n and $\tilde{w}_n$ in the Orlov operators are defined by the above canonical relations and the following flow equations

$$\begin{aligned} \frac{\partial M}{\partial T_n} &= \{B_n, M\}, \qquad & \frac{\partial M}{\partial \tilde{T}_n} &= \{\tilde{B}_n, M\}, \\ \frac{\partial \tilde{M}}{\partial T_n} &= \{B_n, \tilde{M}\}, \qquad & \frac{\partial \tilde{M}}{\partial \tilde{T}_n} &= \{\tilde{B}_n, \tilde{M}\}. \end{aligned} \tag{4.5}$$

Inspired by the twistor construction (or Riemann-Hilbert problem) for the solution structure of the dToda hierarchy [14,11], we now give the twistor construction for the generalized Benney hierarchy.

THEOREM 1 [20]. Let $f(p,X), g(p,X), \tilde{f}(p,X), \tilde{g}(p,X)$ be functions satisfying

$$\{f(p,X), g(p,X)\} = 1, \qquad \{\tilde{f}(p,X), \tilde{g}(p,X)\} = 1. \tag{4.6}$$

Then the functional equations

$$f(\mu, M) = \tilde{f}(\tilde{\mu}, \tilde{M}), \qquad g(\mu, M) = \tilde{g}(\tilde{\mu}, \tilde{M}) \tag{4.7}$$

provide a solution of (4.2) and (4.5). We call the pairs (f, g) and $(\tilde{f}, \tilde{g})$ the twistor data of the solution.

To apply the above Theorem to the present case, we have to impose the following constraint

$$L = \mu^2 = \tilde{\mu}^{-1}, \tag{4.8}$$

that is, $f(p,X) = p^2$ and $\tilde{f}(p,X) = p^{-1}$. As a result, the time variables $\tilde{T}_n$ can be eliminated via the following identification:

$$\tilde{T}_n = -T_{2n}.$$

From (4.6), the twistor data $g(p,X)$ and $\tilde{g}(p,X)$ can be assumed to be in the following form

$$g(p,X) = \frac{X}{2p} - \sum_{n=1}^{\infty} nT_{2n}p^{2n-2}, \qquad \tilde{g}(p,X) = -Xp^2, \tag{4.9}$$

where the second part of $g(p,X)$ is responsible for the string equations (see below). By the Theorem 1 and equation (4.9), we get the following constraint for the Orlov operators:

$$\frac{M\mu^{-1}}{2} - \sum_{n=1}^{\infty} nT_{2n}\mu^{2n-2} = -\tilde{\mu}^2\tilde{M}. \tag{4.10}$$

It's the above constraint that leads to the string equations. So far, the twistor construction only involves the Lax flows (1.2). To obtain the string equations associated with the underlying TFT described in sec. 3, we have to properly extend the standard Orlov operator to include the additional hierarchy equations (2.8). Namely, it's necessary to introduce the flows generated by the logarithmic operator

$$\bar{B}_n = [L^n(\log L - c_n)]_{\geq 1}.$$

Let $\bar{T}_n$ be the time variables of additional flows generated by $\bar{B}_n$ and then the Orlov operator M should be deformed by these new flows to M' so that

$$\frac{\partial M'}{\partial \bar{T}_n} = 2\{\bar{B}_n, M'\}. \tag{4.11}$$

To construct the modified Orlov operator M', it is convenient to using the dressing method [11]. Let us first express the original Lax operator μ and its conjugate Orlov operator M in dressing form

$$\mu = e^{\mathrm{ad}\Theta}(p), \qquad M = e^{\mathrm{ad}\Theta}\left(\sum_{n=1}^{\infty} nT_n p^{n-1}\right) \tag{4.12}$$

where

$$e^{\mathrm{ad}_f}(g) = g + \{f, g\} + \frac{1}{2!}\{f, \{f, g\}\} + \cdots.$$

One can understand that this is the canonical transformation generated by $\Theta(T, \bar{T}, p)$ (for the $\bar{T}$-dependence, see below) and its flow equations (Sato equations) can be written as [11]

$$\nabla_{T_n,\Theta}\Theta = B_n - e^{\mathrm{ad}\Theta}(p^n) \tag{4.13}$$

where

$$\nabla_{T_n,\Theta} f \equiv \sum_{k=0}^{\infty} \frac{1}{(k+1)!}(\mathrm{ad}\Theta)^k \left(\frac{\partial f}{\partial T_n}\right).$$

It is easy to show that (4.13) together with (4.12) implies the flow equations (4.5). In contrast with equation (4.13), the $\bar{T}_n$ flows for Θ are given by

$$\nabla_{\bar{T}_n,\Theta}\Theta = 2\bar{B}_n - 2e^{\mathrm{ad}\Theta}[p^{2n}(\log p^2 - c_n)]$$

and a similar argument reaches the modified Orlov operator

$$\begin{aligned} M' &= e^{\mathrm{ad}\Theta}\left(\sum_{n=1}^{\infty} nT_n p^{n-1} + 4\sum_{n=1}^{\infty} n\bar{T}_n p^{2n-1}(\log p^2 - c_{n-1})\right) \\ &= \sum_{n=1}^{\infty} 2nT_{2n}\mu^{2n-1} + \sum_{n=0}^{\infty}(2n+1)T_{2n+1}\mu^{2n} + \sum_{n=1}^{\infty} w_n \mu^{-n} \\ &\quad + 4\sum_{n=1}^{\infty} n\bar{T}_n \mu^{2n-1}(\log \mu^2 - c_{n-1}), \end{aligned} \tag{4.14}$$

which satisfies the additional flow equations (4.11).

Now we are in the position to derive the string equation. Let us decompose $(M'\mu^{-1}/2 - \sum_{n=1}^{\infty} nT_{2n}\mu^{2n-2})$ into the positive power part $(M'\mu^{-1}/2 - \sum_{n=1}^{\infty} nT_{2n}\mu^{2n-2})_{\geq 1}$ and the non-positive power part $(M'\mu^{-1}/2 - \sum_{n=1}^{\infty} nT_{2n}\mu^{2n-2})_{\leq 0}$ by using equations (4.10) and (4.14). It turns out that

$$\begin{aligned}\left(\frac{M'\mu^{-1}}{2} - \sum_{n=1}^{\infty} nT_{2n}\mu^{2n-2}\right)_{\geq 1} &= \sum_{n=1}^{\infty} \frac{(2n+1)}{2} T_{2n+1}B_{2n-1} + 2\sum_{n=1}^{\infty} n\bar{T}_n\bar{B}_{n-1},\\ \left(\frac{M'\mu^{-1}}{2} - \sum_{n=1}^{\infty} nT_{2n}\mu^{2n-2}\right)_{\leq 0} &= -\left(\tilde{\mu}^2\tilde{M}\right)_{\leq 0} = -\left(\sum_{n=1}^{\infty} nT_{2n}\mu^{2(n-1)}\right)_{\leq 0}\end{aligned}$$

where, by the definition (4.1), we have used the fact that $(\mu^{-2n})_{\geq 1} = (\tilde{\mu}^n)_{\leq 0} = 0$ for $n \geq 1$. Hence

$$g(\mu, M') = \sum_{n=1}^{\infty} \frac{(2n+1)}{2} T_{2n+1}B_{2n-1} + 2\sum_{n=1}^{\infty} n\bar{T}_n\bar{B}_{n-1} - \left(\sum_{n=1}^{\infty} nT_{2n}\mu^{2(n-1)}\right)_{\leq 0}$$

which together with the canonical commutation relation $\{g(\mu, M'), L\} = -1$ implies

$$\sum_{n=2}^{\infty} nT_{2n}\frac{\partial L}{\partial T_{2n-2}} + \sum_{n=1}^{\infty} \frac{(2n+1)}{2} T_{2n+1}\frac{\partial L}{\partial T_{2n-1}} + \sum_{n=1}^{\infty} n\bar{T}_n\frac{\partial L}{\partial \bar{T}_{n-1}} = -1$$

where the Lax flows (1.2) and (2.8) have been used. Therefore by (3.5) we obtain that

$$\sum_{n=1}^{\infty} nT_{R,n}\frac{\partial L}{\partial T^{R,n-1}} + \sum_{n=1}^{\infty} nT^{P,n}\frac{\partial L}{\partial T^{P,n-1}} + \sum_{n=1}^{\infty} nT^{Q,n}\frac{\partial L}{\partial T^{Q,n-1}} = -1.$$

After shifting $T^{P,1} \to T^{P,1} - 1$, we get

$$\sum_{n=1}^{\infty} nT^{R,n}\frac{\partial L}{\partial T^{R,n-1}} + \sum_{n=1}^{\infty} nT^{P,n}\frac{\partial L}{\partial T^{P,n-1}} + \sum_{n=1}^{\infty} nT^{Q,n}\frac{\partial L}{\partial T^{Q,n-1}} + 1 = \frac{\partial L}{\partial T^{P,0}}.$$

Comparing the both-hand sides of the above equation in p and taking into account the constitutive relations (3.2) we obtain the following genus-zero string equation

$$\begin{aligned}t^1(T) &= T^P + 2\sum_{n=1}^{\infty}\sum_{\alpha=P,Q,R} nT^{\alpha,n}\langle\sigma_{n-1}(\mathcal{O}_\alpha)P\rangle,\\ t^2(T) &= T^Q + 2\sum_{n=1}^{\infty}\sum_{\alpha=P,Q,R} nT^{\alpha,n}\langle\sigma_{n-1}(\mathcal{O}_\alpha)R\rangle,\\ t^3(T) &= T^R + 2\sum_{n=1}^{\infty}\sum_{\alpha=P,Q,R} nT^{\alpha,n}\langle\sigma_{n-1}(\mathcal{O}_\alpha)Q\rangle\end{aligned}$$

which gives a special solution of the generalized Benney hierarchy and is used to characterize the free energy in the full phase space. As a consistent check, differentiating t^α to $T^{\beta,n}$ and restricting to the small phase space we recover the flows (3.7) as expected.

V. QUANTIZATION OF THE POISSON BRACKETS

So far, the TFT approach to the generalized Benney hierarchy is restricted to the genus-zero sector. In general it is not easy to promote the Benney system to higher genera through the TFT formulation due to the complexity of topological recursion relations. Nevertheless there are at least two approaches enable us to get consistent genus-one extensions for some integrable hierarchies. One of them is based on the large-N matrix expansion of integrable hierarchies developed by Eguchi and Yang [6] and the other, the theory of Frobenius manifolds by DZ [33]. In this section, we shall compute the genus-one correction to the bi-Hamiltonian structure of the generalized Benney hierarchy by using the latter approach. Roughly speaking, the DZ approach consists of the following two main steps:

- introducing slow spatial and time variables scaling

$$T^{\alpha,n}\to\epsilon T^{\alpha,n},$$

- changing the full free energy as

$$\mathcal{F}\to\sum_{g=0}^{\infty}\epsilon^{2g-2}\mathcal{F}_g,$$

where ϵ is the parameter of genus expansion. As a result, all of the corrections become series in ϵ.

To get a unambiguous genus-one correction of the Hamiltonian flows (3.6) it is convenient to expand the flat coordinates up to the ϵ^2 order as

$$t_\alpha=t_\alpha^{(0)}+\epsilon^2t_\alpha^{(1)}+O(\epsilon^4),\qquad t_\alpha=\eta_{\alpha\beta}t^\beta$$

where $t_\alpha^{(0)}$ is the ordinary flat coordinates t_α satisfying (3.7) and $t_\alpha^{(1)}$ is the genus-one correction defined by $t_\alpha^{(1)}=\partial^2\mathcal{F}_1(T)/\partial T^\alpha\partial X$. Then there exists a unique hierarchy flows of the form [33]

$$\frac{\partial t^\alpha}{\partial T^{\beta,n}}=\{t^\alpha(X),H_{\beta,n+1}\}_1=\{t^\alpha(X),H_{\beta,n}\}_2\tag{5.1}$$

with

$$\begin{aligned}\{t^\alpha(X),t^\beta(Y)\}_i&=\{t^\alpha(X),t^\beta(Y)\}_i^{(0)}+\epsilon^2\{t^\alpha(X),t^\beta(Y)\}_i^{(1)}+O(\epsilon^4),\qquad i=1,2\\ H_{\alpha,n}&=H_{\alpha,n}^{(0)}+\epsilon^2H_{\alpha,n}^{(1)}+O(\epsilon^4)\end{aligned}$$

That means under such correction the Poisson brackets J_1 and J_2 and the Hamiltonians will receive corrections up to ϵ^2 such that the Hamiltonian flows (5.1) still commute with each other. In [33], the genus-one part of the free energy can be expressed as

$$\mathcal{F}_1(T) = \left[\frac{1}{24}\log\det M^\alpha_\beta(t, \partial_X t) + G(t)\right]_{t=t(T)}, \tag{5.2}$$

where the matrix M^α_β is given by

$$M^\alpha_\beta(t, \partial_X t) = c^\alpha_{\beta\gamma}(t)\partial_X t^\gamma = \begin{pmatrix} t^1_X & t^3_X & t^2_X \\ t^2_X & t^1_X + \frac{t^3 t^3_X}{2} & \frac{t^3 t^2_X + t^3_X t^2}{2} \\ t^3_X & \frac{2t^2_X}{t^2} & t^1_X + \frac{t^3 t^3_X}{2} \end{pmatrix}$$

and $G(t^2, t^3)$ is the associated G-function satisfying the Getzler's equation (1.7) which are over-determined and some of them are redundant. In general, they can be solved explicitly using canonical coordinates for an arbitrary semi-simple Frobenius manifold [29]. Here we notice that $c^3_{22} = 2/t^2$, and $c^2_{22} = c^1_{22} = 0$ and substitute them into the coefficient of $(z_2)^4$ of the Getzler equations (1.7) to get the following ordinary differential equation

$$\frac{\partial^2 G}{\partial (t^2)^2} + \frac{1}{2t^2}\frac{\partial G}{\partial t^2} - \frac{1}{24(t^2)^2} = 0$$

which together with the quasi-homogeneous property $\mathcal{L}_E G = -1/8$ (c.f. Theorem 3 in [33]) implies (up to a constant)

$$G(t_2, t_3) = -\frac{\log t^2}{12}. \tag{5.3}$$

We remark that the above G-function is the same as the one of the Benney hierarchy with two primary fields [20].

Thus a simple computation yields

$$\mathcal{F}_1 = \frac{1}{24}\log\left[t^2(t^1_X)^3 + 2(t^2_X)^3 + \frac{(t^2)^2(t^3_X)^3}{2} + t^2 t^3 (t^1_X)^2 t^3_X + \frac{t^2(t^3_X)^2(t^3)^2 t^2 t^1_X}{4}\right.$$
$$\left. - t^3 t^1_X (t^2_X)^2 - 3t^2 t^1_X t^2_X t^3_X - \frac{t^2 t^3 t^2_X (t^3_X)^2}{2}\right] - \frac{\log t^2}{8}.$$

Using $c^\alpha_{\beta\gamma}$ and $\mathcal{F}_1$ and consulting the procedure developed by DZ (c.f Theorems 1 and 2 in [33]) it turns out that the genus-one corrections of the Poisson brackets, in terms of flat coordinates t^α, are given by

$$\{t^1(X), t^2(Y)\}_1 = \{t^2(X), t^2(Y)\}_1 = \{t^3(X), t^3(Y)\}_1 = O(\epsilon^4),$$
$$\{t^1(X), t^1(Y)\}_1 = 2\delta'(X-Y) + O(\epsilon^4),$$
$$\{t^1(X), t^3(Y)\}_1 = -\frac{2\epsilon^2}{3}\left[\frac{\delta'''(X-Y)}{t^2} - \frac{2t^2_X \delta''(X-Y)}{(t^2)^2} + \frac{2(t^2_X)^2\delta'(X-Y)}{(t^2)^3}\right.$$

$$-\frac{t^2_{XX}\delta'(X-Y)}{(t^2)^2}\Bigg] + O(\epsilon^4),$$

$$\{t^2(X), t^3(Y)\}_1 = 2\delta'(X-Y) - \frac{\epsilon^2}{3}\left[\frac{t^3\delta'''(X-Y)}{t^2} - \frac{2t^3t^2_X\delta''(X-Y)}{(t^2)^2} + \frac{t^3_X\delta''(X-Y)}{t^2}\right.$$
$$\left. -\frac{t^3_Xt^2_X\delta'(X-Y)}{(t^2)^2} - \frac{t^3t^2_{XX}\delta'(X-Y)}{(t^2)^2} + \frac{2t^3(t^2_X)^2\delta'(X-Y)}{(t^2)^3}\right] + O(\epsilon^4),$$

and

$$\{t^1(X), t^1(Y)\}_2 = t^1_X\delta(X-Y) + 2t^1\delta'(X-Y) + \frac{\epsilon^2\delta'''(X-Y)}{2} + O(\epsilon^4),$$
$$\{t^1(X), t^2(Y)\}_2 = 2t^2_X\delta(X-Y) + 3t^2\delta'(X-Y) + O(\epsilon^4),$$
$$\{t^1(X), t^3(Y)\}_2 = t^3\delta'(X-Y) + \epsilon^2\left[-\frac{4t^1+(t^3)^2}{6t^2}\delta'''(X-Y)\right.$$
$$+\frac{4t^1t^2_X + (t^3)^2t^2_X - 4t^2t^1_X - 2t^2t^3t^3_X}{3(t^2)^2}\delta''(X-Y)$$
$$\left.+\left(\frac{4t^1t^2_X + (t^3)^2t^2_X - 4t^2t^1_X - 2t^2t^3t^3_X}{6(t^2)^2}\right)_X\delta'(X-Y)\right] + O(\epsilon^4),$$
$$\{t^2(X), t^2(Y)\}_2 = 2t^2t^3\delta'(X-Y) + (t^2t^3)_X\delta(X-Y)$$
$$+\epsilon^2\left[\frac{1}{3}\left(2t^1_{XX} + (t^3_X)^2 + t^3t^3_{XX} - \frac{2t^2_Xt^1_X}{t^2} - \frac{t^2_Xt^3t^3_X}{t^2}\right)\delta'(X-Y)\right.$$
$$\left.+\frac{1}{6}\left(2t^1_{XXX} + 3t^3_Xt^3_{XX} + t^3t^3_{XXX} - \left(\frac{t^3t^3_Xt^2_X + 2t^1_Xt^2_X}{t^2}\right)_X\right)\delta(X-Y)\right] + O(\epsilon^4),$$
$$\{t^2(X), t^3(Y)\}_2 = \left(2t^1 + \frac{(t^3)^2}{2}\right)\delta'(X-Y) + \epsilon^2\left[\left(1 - \frac{t^1t^3}{3t^2} - \frac{(t^3)^3}{12t^2}\right)\delta'''(X-Y)\right.$$
$$+\left(\frac{8t^3t^1t^2_X + 2(t^3)^3t^2_X - 8t^2t^3t^1_X - 5t^2(t^3)^2t^3_X - 4t^2t^1t^3_X - 4t^2t^2_X}{12(t^2)^2}\right)\delta''(X-Y)$$
$$+\left(\left(\frac{t^3t^2_X}{3(t^2)^2}\right)_X t^1 - \frac{(t^3)^3(t^2_X)^2}{6(t^2)^3} + \frac{5(t^3)^2t^2_Xt^3_X}{12(t^2)^2} - \frac{t^3(t^3_X)^2}{3t^2} - \frac{(t^3)^2t^3_{XX}}{6t^2}\right.$$
$$\left.\left.-\frac{t^3_Xt^1_X}{3t^2} + \frac{(t^3)^3t^2_{XX}}{12(t^2)^2} + \frac{2t^3t^2_Xt^1_X}{3(t^2)^2} - \frac{(t^2_X)^2}{6(t^2)^2} - \frac{t^3t^1_{XX}}{3t^2}\right)\delta'(X-Y)\right] + O(\epsilon^4),$$
$$\{t^3(X), t^3(Y)\}_2 = 6\delta'(X-Y) - \epsilon^2\left[\frac{2t^3}{3t^2}\delta'''(X-Y) + \left(\frac{t^3}{t^2}\right)_X\delta''(X-Y)\right.$$
$$\left.+\left(\frac{t^3}{3t^2}\right)_{XX}\delta'(X-Y)\right] + O(\epsilon^4)$$

which still satisfy the relation $J_1^{ij} = \partial J_2^{ij}/\partial t^1$. For the genus-one corrections to the Hamiltonians $H_{\alpha,n}$ we collect some of them in Appendex B.

VI. CONCLUDING REMARKS

We have studied the generalized Benney hierarchy from the topological field theory . First of all, we can construct the free energy from the bi-Hamiltonian structure of the generalized Benney hierarchy using the flat pencil. Secondly, after introducing the flat coordinates, the primary fields in the LG formulation of TFT can be defined. Then the topological recursion relations between gravitational descendants at genus-zero turn out to be the associated commuting Hamiltonian flows in the generalized Benney hierarchy. Furthermore, we use the twistor construction to obtain the string equation associated with the TFT, which is a Galiean-type symmetry of the generalized Benney hierarchy and characterizes the free energy in the full phase space. Finally, based on the approach of DZ, we obtain the genus-one corrections of the Poisson brackets.

In spite of the results obtained, there are some interesting issues deserving more investigations:

- In [20], it can be shown that the string coupling constant ϵ of the Benney hierarchy will correspond to the effect of dispersion of Kaup-Broer hierarchy [36,37]. However, for the case of the generalized Benney hierarchy, the computation is more involved and needs further investigations.
- The relations of free energies between dKP and dmKP are interesting. In [15], the free energy of dKP is constructed using generalized Gelfand-Dickey potential without referring to the bi-Hamiltonian structure of dKP. A similar approach for dmKP should be developed and see what role is played by the Miura map between dKP and dmKP [18].
- Recently, the associativity equations of infinite many primary fields in dKP are obtained [38] by the dispersionless Hirota equations of dKP [39]. Thus it is quite natural to ask whether the dmKP hierarchy and its reductions have the similar structures of associativity equations and what kind of infinite dimensional Frobenius manifold [40] they provide.

Acknowledgements

JHC thanks Y. Kodama for helpful discussions. MHT thanks the National Science Council of Taiwan (Grant No. NSC 89-2112-M194-020) for support.

APPENDIX A: PRIMARY FLOWS AND TWO-POINT FUNCTIONS

1. Primary $T^{P,n}$-flows

$$
\begin{aligned}
\frac{1}{2}\frac{\partial v^1}{\partial T^{P,0}} &= \langle PQ\rangle_X,\\
\frac{1}{2}\frac{\partial v^2}{\partial T^{P,0}} &= \langle QR\rangle_X,\\
\frac{1}{2}\frac{\partial v^3}{\partial T^{P,0}} &= \langle PR\rangle_X,\\
\frac{1}{2}\frac{\partial v^1}{\partial T^{P,1}} &= \langle \sigma_1(P)PQ\rangle = \left[\frac{1}{2}v^1v^2 - \frac{1}{24}(v^1)^3 + v^3\right]_X,\\
\frac{1}{2}\frac{\partial v^2}{\partial T^{P,1}} &= \langle \sigma_1(P)QR\rangle = \frac{1}{2}v^2v^2_X + \frac{1}{8}(v^1)^2v^2_X + v^1v^3_X + \frac{1}{2}v^1_Xv^3,\\
\frac{1}{2}\frac{\partial v^3}{\partial T^{P,1}} &= \langle \sigma_1(P)PR\rangle = \left[\frac{1}{2}v^2v^3 + \frac{1}{8}(v^1)^2v^3\right]_X,\\
\frac{1}{2}\frac{\partial v^1}{\partial T^{P,2}} &= \langle \sigma_2(P)PQ\rangle = \left[\frac{1}{160}(v^1)^5 - \frac{1}{12}(v^1)^3v^2 + \frac{1}{2}(v^1)^2v^3 + \frac{1}{2}v^1(v^2)^2 + 2v^2v^3\right]_X,\\
\frac{1}{2}\frac{\partial v^2}{\partial T^{P,2}} &= \langle \sigma_2(P)QR\rangle = -\frac{1}{96}(v^1)^4v^2_X + \frac{1}{6}(v^1)^3v^3_X + \frac{1}{4}(v^1)^2v^2v^2_X + \frac{1}{4}(v^1)^2v^1_Xv^3 + 2v^1(v^2v^3)_X\\
&\quad + \frac{1}{6}(v^2)^3_X + (v^3)^2_X + v^1_Xv^2v^3,\\
\frac{1}{2}\frac{\partial v^3}{\partial T^{P,2}} &= \langle \sigma_2(P)PR\rangle = \left[-\frac{1}{96}(v^1)^4v^3 + \frac{1}{4}(v^1)^2v^2v^3 + v^1(v^3)^2 + \frac{1}{2}(v^2)^2v^3\right]_X.
\end{aligned}
$$

From above, we can extract the following two-point functions:

$$
\begin{aligned}
\langle \sigma_1(P)P\rangle &= \frac{1}{64}(v^1)^4 - \frac{1}{8}(v^1)^2v^2 + \frac{1}{2}v^1v^3 + \frac{1}{4}(v^2)^2,\\
\langle \sigma_1(P)Q\rangle &= \frac{1}{2}v^1v^2 - \frac{1}{24}(v^1)^3 + v^3,\\
\langle \sigma_1(P)R\rangle &= \frac{1}{2}v^2v^3 + \frac{1}{8}(v^1)^2v^3,\\
\langle \sigma_2(P)P\rangle &= -\frac{1}{384}(v^1)^6 + \frac{1}{32}(v^1)^4v^2 - \frac{1}{12}(v^1)^3v^3 - \frac{1}{8}(v^1)^2(v^2)^2 + v^1v^2v^3 + \frac{1}{6}(v^2)^3 + (v^3)^2,\\
\langle \sigma_2(P)Q\rangle &= \frac{1}{160}(v^1)^5 - \frac{1}{12}(v^1)^3v^2 + \frac{1}{2}(v^1)^2v^3 + \frac{1}{2}v^1(v^2)^2 + 2v^2v^3,\\
\langle \sigma_2(P)R\rangle &= -\frac{1}{96}(v^1)^4v^3 + \frac{1}{4}(v^1)^2v^2v^3 + v^1(v^3)^2 + \frac{1}{2}(v^2)^2v^3.
\end{aligned}
$$

2. Primary $T^{Q,n}$-flows

$$\begin{aligned}
\frac{1}{2}\frac{\partial v^1}{\partial T^{Q,0}} &= (\log(v^3))_X,\\
\frac{1}{2}\frac{\partial v^2}{\partial T^{Q,0}} &= \frac{1}{2}v^1_X + \frac{1}{2}v^1(\log(v^3))_X,\\
\frac{1}{2}\frac{\partial v^3}{\partial T^{Q,0}} &= \frac{1}{2}v^2_X,\\
\frac{1}{2}\frac{\partial v^1}{\partial T^{Q,1}} &= \langle\sigma_1(Q)PQ\rangle = \left[\frac{1}{4}(v^1)^2 + v^2\log v^3\right]_X,\\
\frac{1}{2}\frac{\partial v^2}{\partial T^{Q,1}} &= \langle\sigma_1(Q)QR\rangle = \frac{1}{2}(v^1v^2)_X + v^3_X\log v^3 + \frac{1}{2}v^1v^2(\log v^3)_X + \frac{1}{2}v^1v^2_X\log v^3,\\
\frac{1}{2}\frac{\partial v^3}{\partial T^{Q,1}} &= \langle\sigma_1(Q)PR\rangle = \left[\frac{1}{4}(v^2)^2 + \frac{1}{2}v^1v^3\log v^3\right]_X,\\
\frac{1}{2}\frac{\partial v^1}{\partial T^{Q,2}} &= \langle\sigma_2(Q)PQ\rangle = \left[-\frac{1}{24}(v^1)^4 + \frac{1}{2}v^2(v^1)^2 + (v^2)^2\log v^3 + 2v^1v^3(\log v^3 - 1)\right]_X,\\
\frac{1}{2}\frac{\partial v^2}{\partial T^{Q,2}} &= \langle\sigma_2(Q)QR\rangle = \frac{1}{2}(v^1(v^2)^2)_X - \frac{1}{6}(v^2(v^1)^3)_X + 2(v^2v^3(\log v^3 - 1))_X + v^1_Xv^1v^3\log v^3\\
&\qquad + (v^1)^2v^3_X\log v^3 + \frac{1}{2}(\frac{1}{2}(v^1)^2v^2 + (v^2)^2\log v^3)_X,\\
\frac{1}{2}\frac{\partial v^3}{\partial T^{Q,2}} &= \langle\sigma_2(Q)PR\rangle = \left[\frac{1}{6}(v^2)^3 + \frac{1}{12}(v^1)^3v^3 + v^1v^2v^3\log v^3 + (v^3)^2(\log v^3 - 2)\right]_X.
\end{aligned}$$

From above, we can extract the following two-point functions:

$$\begin{aligned}
\langle\sigma_1(Q)P\rangle &= \frac{1}{2}v^1v^2 - \frac{1}{12}(v^1)^3 + v^3(\log v^3 - 1),\\
\langle\sigma_1(Q)Q\rangle &= \frac{1}{4}(v^1)^2 + v^2\log v^3,\\
\langle\sigma_1(Q)R\rangle &= \frac{1}{4}(v^2)^2 + \frac{1}{2}v^1v^3\log v^3,\\
\langle\sigma_2(Q)P\rangle &= \frac{1}{60}(v^1)^5 - \frac{1}{6}(v^1)^3v^2 + \frac{1}{2}(v^1)^2v^3 + \frac{1}{2}v^1(v^2)^2 + 2v^2v^3(\log v^3 - 1),\\
\langle\sigma_2(Q)Q\rangle &= -\frac{1}{24}(v^1)^4 + \frac{1}{2}v^2(v^1)^2 + (v^2)^2\log v^3 + 2v^1v^3(\log v^3 - 1),\\
\langle\sigma_2(Q)R\rangle &= \frac{1}{6}(v^2)^3 + \frac{1}{12}(v^1)^3v^3 + v^1v^2v^3\log v^3 + (v^3)^2(\log v^3 - 2).
\end{aligned}$$

3. Primary $T^{R,n}$-flows

$$
\begin{aligned}
\frac{1}{2}\frac{\partial v^1}{\partial T^{R,0}} &= \frac{1}{2}v^2_X,\\
\frac{1}{2}\frac{\partial v^2}{\partial T^{R,0}} &= (\frac{1}{4}v^1v^2_X + \frac{1}{2}v^3_X),\\
\frac{1}{2}\frac{\partial v^3}{\partial T^{R,0}} &= \frac{1}{4}(v^1v^3)_X,\\
\frac{1}{2}\frac{\partial v^1}{\partial T^{R,1}} &= \langle\sigma_1(R)PQ\rangle = \left[\frac{1}{2}v^1v^3 + \frac{1}{4}(v^2)^2\right]_X,\\
\frac{1}{2}\frac{\partial v^2}{\partial T^{R,1}} &= \langle\sigma_1(R)QR\rangle = \frac{1}{2}(v^3v^2)_X + \frac{1}{4}(v^1v^1_Xv^3 + v^1v^2_Xv^2 + (v^1)^2v^3_X),\\
\frac{1}{2}\frac{\partial v^3}{\partial T^{R,1}} &= \langle\sigma_1(R)PR\rangle = \left[\frac{1}{4}(v^3)^2 + \frac{1}{4}v^1v^2v^3\right]_X,\\
\frac{1}{2}\frac{\partial v^1}{\partial T^{R,2}} &= \langle\sigma_2(R)PQ\rangle = \left[\frac{1}{6}(v^2)^3 + v^2v^1v^3 + \frac{1}{2}(v^3)^2\right]_X,\\
\frac{1}{2}\frac{\partial v^2}{\partial T^{R,2}} &= \langle\sigma_2(R)QR\rangle = \frac{1}{2}((v^2)^2v^3)_X + \frac{1}{2}(v^1)^2(v^2v^3)_X + \frac{1}{2}v^1v^1_Xv^2v^3 + \frac{3}{2}v^1v^3v^3_X\\
&\quad +\frac{1}{2}v^1_X(v^3)^2 + \frac{1}{4}v^1(v^2)^2v^2_X,\\
\frac{1}{2}\frac{\partial v^3}{\partial T^{R,2}} &= \langle\sigma_2(R)PR\rangle = \left[\frac{1}{4}(v^1)^2(v^3)^2 + \frac{1}{4}v^1(v^2)^2v^3 + \frac{1}{2}v^2(v^3)^2\right]_X.
\end{aligned}
$$

From above, we can extract the following two-point functions:

$$
\begin{aligned}
\langle\sigma_1(R)P\rangle &= \frac{1}{2}v^2v^3,\\
\langle\sigma_1(R)Q\rangle &= \frac{1}{2}v^1v^3 + \frac{1}{4}(v^2)^2,\\
\langle\sigma_1(R)R\rangle &= \frac{1}{4}(v^3)^2 + \frac{1}{4}v^1v^2v^3,\\
\langle\sigma_2(R)P\rangle &= \frac{1}{2}(v^2)^2v^3 + \frac{1}{2}v^1(v^3)^2,\\
\langle\sigma_2(R)Q\rangle &= \frac{1}{6}(v^2)^3 + v^2v^1v^3 + \frac{1}{2}(v^3)^2,\\
\langle\sigma_2(R)R\rangle &= \frac{1}{4}(v^1)^2(v^3)^2 + \frac{1}{4}v^1(v^2)^2v^3 + \frac{1}{2}v^2(v^3)^2.
\end{aligned}
$$

APPENDIX B: GENUS-ONE CORRECTIONS FOR THE HAMILTONIANS

$$H_{P,0} = \int \frac{t^1}{2} + O(\epsilon^4),$$

$$
\begin{aligned}
H_{P,1} &= \int \left[\frac{1}{2}t^2t^3 + \frac{1}{4}(t^1)^2\right] - \epsilon^2 \int \left[\frac{t^1_X t^2_X}{6t^2} + \frac{t^2_X t^3 t^3_X}{12t^2}\right] + O(\epsilon^4),\\
H_{P,2} &= \frac{1}{6}\int \left[\frac{1}{2}(t^1)^3 + \frac{1}{2}t^2(t^3)^3 + 3t^1t^2t^3 + 3(t^2)^2\right] - \epsilon^2 \int \biggl[\frac{5}{24}t^1_X t^1_X + \frac{t^3 t^2_X t^2_X}{4t^2} + \frac{(t^3)^2 t^3_X t^3_X}{12}\\
&\qquad + \frac{((t^3)^2 + 2t^1)t^1_X t^2_X}{12t^2} + \frac{t^3 t^1_X t^3_X}{4} + \left(\frac{1}{3} + \frac{t^1t^3}{12t^2} + \frac{(t^3)^3}{24t^2}\right) t^2_X t^3_X\biggr] + O(\epsilon^4),\\
H_{R,0} &= \int \frac{t^2}{2} + O(\epsilon^4),\\
H_{R,1} &= \frac{1}{2}\int \left[t^1t^2 + \frac{t^2(t^3)^2}{4}\right] - \frac{\epsilon^2}{4}\int \left[\frac{t^3(t^3_X)^2}{6} + \frac{t^3 t^2_X t^1_X}{3t^2} + \frac{t^2_X t^3_X (t^3)^2}{6t^2} + \frac{(t^2_X)^2}{2t^2} + \frac{t^3_X t^1_X}{3}\right] + O(\epsilon^4),\\
H_{R,2} &= \frac{1}{4}\int \left[t^3(t^2)^2 + t^2(t^1)^2 + \frac{1}{2}t^2t^1(t^3)^2 + \frac{1}{16}t^2(t^3)^4\right] - \frac{\epsilon^2}{4}\int \biggl[\frac{t^3}{3}t^1_X t^1_X + \left(\frac{t^1}{2t^2} + \frac{3(t^3)^2}{8t^2}\right) t^2_X t^2_X\\
&\qquad + \left(\frac{t^2}{4} + \frac{t^1t^3}{6} + \frac{(t^3)^3}{8}\right) t^3_X t^3_X + \left(\frac{5}{3} + \frac{t^1t^3}{3t^2} + \frac{(t^3)^3}{12t^2}\right) t^1_X t^2_X + \left(\frac{t^1}{3} + \frac{5(t^3)^2}{12}\right) t^1_X t^3_X\\
&\qquad + \left(\frac{4t^3}{3} + \frac{t^1(t^3)^2}{6t^2} + \frac{(t^3)^4}{24t^2}\right) t^2_X t^3_X\biggr] + O(\epsilon^4),\\
H_{Q,0} &= \int \frac{t^3}{2} + O(\epsilon^4),\\
H_{Q,1} &= \int \left[\frac{1}{24}(t^3)^3 + \frac{1}{2}t^1t^3 + t^2\left(\log t^2 - \frac{5}{2}\right)\right] - \epsilon^2 \int \biggl[\frac{1}{12t^2}t^1_X t^1_X + \frac{(t^3)^2(t^3_X)^2}{48t^2}\\
&\qquad + \frac{t^2_X t^3_X}{12t^2} + \frac{(t^2_X)^2 t^3}{24(t^2)^2} + \frac{t^3 t^3_X t^1_X}{12t^2}\biggr] + O(\epsilon^4),\\
H_{Q,2} &= \int \left[\frac{(t^3)^5}{320} + \frac{t^1(t^3)^3}{24} + \frac{(t^1)^2t^3}{4} + \frac{1}{4}(t^3)^2t^2\left(\log t^2 - \frac{3}{2}\right) + t^1t^2\left(\log t^2 - \frac{5}{2}\right)\right]\\
&\qquad -\epsilon^2 \int \biggl[\frac{1}{6t^2}\left(\frac{1}{8}(t^3)^2 + \frac{t^1}{2}\right) t^1_X t^1_X + \frac{1}{4t^2}\left(\log t^2 + \frac{t^1t^3}{6t^2} + \frac{(t^3)^3}{24t^2} - \frac{3}{2}\right) t^2_X t^2_X\\
&\qquad + \frac{t^3}{12}\left(\log t^2 + \frac{(t^3)^3}{16t^2} + \frac{t^1t^3}{4t^2} - \frac{5}{4}\right) t^3_X t^3_X + \frac{t^3}{6t^2}\left(\log t^2 - \frac{1}{2}\right) t^1_X t^2_X\\
&\qquad + \frac{1}{6}\left(\log t^2 + \frac{(t^3)^3}{8t^2} + \frac{t^1t^3}{2t^2} - 1\right) t^1_X t^3_X + \frac{1}{12t^2}\left(\frac{-3(t^3)^2}{4} + t^1 + (t^3)^2\log t^2\right) t^2_X t^3_X\biggr] + O(\epsilon^4).
\end{aligned}
$$

REFERENCES

[1] R. Dijkgraaf, New Symmetry Principles in Quantum Field Theory, ed. by J. Frohlich *et. al* (New York: Plenum, 1993) pp95-158 and references therein.

[2] E. Witten, Nucl. Phys. B 340 (1990) 281; Surv. Diff. Geom. **1** (1991) 243.

[3] M. Kontsevich, Commun. Math. Phys. 147(1992) 1.

[4] L.A. Dickey, Soliton Equations and Hamiltonian Systems (Singapore: World Scientific, 1991).

[5] I. Krichever, Commun. Math. Phys. 143 (1992) 415; Commun. Pure Appl. Math. 47 (1994) 437.

[6] T. Eguchi and S.K. Yang, Mod. Phys. Lett. A 9 (1994) 2893.

[7] T. Eguchi, K. Hori and S.K. Yang, Int. J. Mod. Phys. A 10 (1995) 4203.

[8] T. Eguchi, Y. Yamada and S.K. Yang, Rev. Math. Phys. 7 (1995) 279.

[9] R. Dijkgraaf, E. Verlinde and H. Verlinde, Nucl. Phys. B 352 (1991) 59.

[10] R. Dijkgraaf and E. Witten, Nucl. Phys. B 342 (1990) 486.

[11] K. Takasaki and T. Takebe, Rev. Math. Phys. 7 (1995) 743.

[12] D. Lebedev and Yu Manin Phys. Lett. A 74 (1979) 154.

[13] Y. Kodama and J. Gibbons, Proceedinds Fourth Workshop on Nonlinear and Turbulent Process in Physics, (Singapore: World Scientific, 1990) p.166.

[14] K. Takasaki and T. Takebe, Lett. Math. Phys. 23 (1991) 205; Int. J. Mod. Phys. A 7 (1992) 889.

[15] S. Aoyama and Y. Kodama, Mod. Phys. Lett. A 9 (1994) 2481; Commun. Math. Phys. 182 (1996) 185.

[16] A. Losev and I. Polyubin, Int. J. Mod. Phys. A10 (1995) 4161.

[17] L.C. Li, Commun. Math. Phys. 203 (1999) 573.

[18] J.H. Chang and M.H. Tu, J. Math. Phys. 41 (2000) 5391.

[19] J.H. Chang and M.H. Tu, J. Math. Phys. 41 (2000) 8117.

[20] J.H. Chang and M.H. Tu, J. Phys. A 34 (2001) 251.

[21] D.J. Benney, Stud. Appl. Math. 52 (1973) 45.

[22] B. Kupershmidt and Yu Manin, Funct. Anal. Appl. 11 (1977) 31 ; ibid. 12 (1978) 25.

[23] V.E. Zakharov, Funct. Anal. Appl. 14 (1980) 89; Physica D 3 (1981) 193.

[24] B. Enriquez, A. Yu Orlov and V.N. Rubtsov, Inverse Problems 12 (1996) 241.

[25] D.B. Fairlie and I.A.B. Strachan, Inverse Problems 12 (1996) 885.

[26] J. Gibbons and S.P. Tsarev, Phys. Lett. A 211 (1996) 19.

[27] B. Dubrovin and S.P. Novikov, Russ. Math. Surv. 44 (1989) 35.

[28] B. Dubrovin, Nucl. Phys. B 379 (1992) 627.

[29] B. Dubrovin, Integrable systems and Quantum Group, ed. by M. Francaviglia and S. Greco (Berlin: Springer, 1996) p.120.

[30] B. Dubrovin, Integrable systems and algebraic geometry, ed. by M.H. Saito, Y. Shimizu and K. Ueno (Singapore: World Scientific, 1998) p.47.

[31] I.A.B. Strachan, J. Math. Phys. 40 (1999) 5058.

[32] C. Vafa, Mod. Phys. Lett. A 6 (1991) 337.

[33] B. Dubrovin and Y. Zhang, Commun. Math. Phys. 198 (1998) 311.
[34] E. Getzler, J. Amer. math. Soc. 10 (1997) 973.
[35] H. Kanno and Y. Ohta, Nucl. Phys. B 442 (1995) 179.
[36] W. Oevel and W. Strampp, Commun. Math. Phys. 157 (1993) 51.
[37] B. Konopelchenko and W. Oevel, Publ. RIMS, Kyoto Univ. 29 (1993) 581.
[38] A. Boyarsky, A. Marshakov, O. Ruchayskiy, P. Wiegmann, A. Zabrodin, "On Associativity Equations in Dispersionless Integrable Hierarchies", hep-th/0105260.
[39] R. Carroll and Y. Kodama, J. Phys. A28 (1995) 6373.
[40] P. Guha and K. Takasaki, J. Geom. Phys. 25 (1998) 326.